

NEW!

IPS e.max® ZirCAD® Prime Esthetic

Yuki Momma, RDT, Boston, MA
"I make it e.max ZirCAD Prime Esthetic
because it provides an elegance
which is unmatched by any other.
It is a game changer!"

For optimal strength, esthetics and multiple indications – Whether you choose lithium disilicate, or zirconia for you next case – Make it e.max!



Learn more about Gradient Technology

ivoclar.com

Making People Smile

ivoclar





Berlin | Chicago | Tokyo Barcelona | London | Milan | Mexico City | Paris | Prague | Seoul | Warsaw Beijing | Istanbul | Sao Paulo | Zagreb



One book, one tree: In support of reforestation worldwide and to address the climate crisis, for every book sold Quintessence Publishing will plant a tree (https://onetreeplanted.org/).

#### Library of Congress Control Number: 2022943698

A CIP record for this book is available from the British Library. ISSN: 1060-1341 / ISBN: 978-1-64724-143-8

#### **PUBLISHER**

Christian W. Haase

#### **PUBLISHER EMERITUS**

H. W. Haase

#### **EXECUTIVE VICE PRESIDENT, DIRECTOR**

William G. Hartman

#### **EDITORIAL**

Leah Huffman

#### **DESIGN/PRODUCTION**

Angelina Schmelter

#### **MANUSCRIPT SUBMISSION**

QDT publishes original articles covering dental laboratory techniques and methods. For QDT submission information, contact the publisher (service@quintbook.com).

Copyright © 2023 by Quintessence Publishing Co, Inc. All rights reserved. No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or any information and retrieval system, without permission in writing from the publisher. The publisher assumes no responsibility for unsolicited manuscripts. All opinions are those of the authors

Permission to photocopy items solely for internal or personal use and for the internal or personal use of specific clients is granted by Quintessence Publishing Co, Inc, for libraries and other users registered with the Copyright Clearance Center (CCC) provided the appropriate fee is paid directly to CCC (www.copyright.com).



© 2023 Quintessence Publishing Co, Inc

Ouintessence Publishing Co, Inc 411 N Raddant Road Batavia, IL 60510 www.quintpub.com

5 4 3 2 1

ISSN: 1060-1341 / ISBN: 978-1-64724-143-8

Printed in Croatia

Cover image courtesy of Joshua Polansky, BA, MDC

# et the Ne Editor-in-Chi

QDT 2023 is the inaugural issue with Master Dental Technician Vincent Fehmer at the helm. Under his editorship, QDT maintains its signature look but with a more fluid design and larger images to showcase the beauty of the work. Read on to learn more about our new editor-in-chief. We are excited to see where QDT goes under his direction!

incent Fehmer received his dental technical education and degree in Stuttgart, Germany, before completing fellowships in Great Britain and the United States in Oral Designcertified dental technical laboratories. After working

several years in such a laboratory in Berlin, he received his MDT degree and became the chief dental technician at the Clinic for Fixed and Removable Prosthodontics in Zurich, Switzerland. Since 2015, he has been a dental technician at the Clinic for Fixed Prosthodontics and Biomaterials in Geneva, Switzerland. He

also runs his own laboratory in Lausanne.

Mr Fehmer is a Fellow of the International Team for Implantology, an active member of the European Academy of Esthetic Dentistry (EAED), and a member of the Oral Design group as well as the European Association of Dental Technology (EADT) and the German Society of Esthetic Dentistry (DGÄZ). He is a sought-after international speaker and has received many honors for his work, including the Kenneth Rudd Award from the American Prosthodontic

> Society. He has published more than 50 articles in peer-reviewed journals within the field of fixed prosthodontics and digital dental technology, contributed to many books, and recently coauthored the book Fixed Restorations: A Clinical Guide to the Selection of Materials and Fabrication Technology with Irena



Vincent Fehmer, MDT, on the lecture circuit.

Sailer and Bjarni Pjetursson. Mr Fehmer also serves as Editor-in-Chief for the International Journal of Esthetic Dentistry, Section Editor for the International Journal of Prosthodontics, and Co-Chair for the 26th International Symposium on Ceramics (June 2023).

# 

EDITORIAL Simplicity Meets Esthetics	viii
Vincent Fehmer	
Enhancing Anterior Esthetics: A Patchwork of Different Types of Ceramics Carola Wohlgenannt	2
Predictable Anterior Esthetics in a Complex Case of a Young Patient with a Congenital Missing Central Incisor Arndt Happe, Anja Krüssmann, and Leonie Debring	22
Laboratory-Generated Digital Concepts in Modern Dental Communication for the Restorative Team: Creating the Virtual Patient Joshua Polansky and Graham Forbes	40
Is Our Job a Lifelong Journey? What Is the Motivation? Is There a Key to Success? Katrin Rohde	53
Facially Driven Full-Mouth Rehabilitation: An Interdisciplinary Approach Mirela Feraru, Ion Nicolescu, Mihai Dorobantu, Michael Tulchinsky, Stefano Inglese, and Nitzan Bichacho	68
The Digital Roadmap for Edentulous Patients in Implant Prosthodontics  Luc Rutten and Patrick Rutten	84

## QUINTESSENCE OF DENTAL TECHNOLOGY

### Volume 45

Central Incisor Mock-up: A Fundamental Role in Esthetic Rehabilitation	116
Giuseppe Romeo, Georgi Iliev, and Galip Gürel	
Interdisciplinary Treatments Using Digital Tools for Diagnosis, Communication, and Execution Gustavo Giordani, Florin Cofar, Eric Van Dooren, Marcelo Giordani, Ioan Barbur, and Ioana Popp	131
Facially Generated Esthetic Planning Shogo Yamamoto and Takuya Yada	148
The Injection Resin Technique Using Digital Design: An Interdisciplinary Approach Douglas A. Terry, Wesam Salha, Memo Castro, and Markus B. Blatz	164
Digital and Analog Approach for Precise and Predictable Restorative Treatments Jun Iwata and Shinya Kuramoto	180
Mucogingival Rotational Tooth Preparation in the Management of Soft Tissues in Fixed Prostheses Ivan Contreras Molina	198
Classification and Indication of Zirconia Restorations According to Layering Design and Thickness Naoki Hayashi	216
Translucency Levels: Full Layering vs Microlayering Aiham Farah, The Aurum Group, and Volinder S. Dhesi	228

# "It is an honor to continue the legacy of such a unique publication..."

#### **EDITORIAL: SIMPLICITY MEETS ESTHETICS**



s the new editor-in-chief of QDT, I am following the vision and work of the outstanding previous editors—Prof John A. Sorensen, Prof Avishai Sadan, and Prof Sillas Duarte—who developed QDT over the last 44 years into a one-of-a-kind, cutting-edge annual publication. Like them, my aim is to connect the leading experts in the field of restorative dentistry and dental technology to elevate this unique publication to the highest standard in our profession.

Important to *my* vision for QDT is the feasibility of the presented work and its actual application to daily practice. My hope is that the articles in these pages inspire clinicians and technicians to apply the techniques they see here for the benefit of all their patients. Therefore, the motto "simplicity meets esthetics" is more than just a connection of words; I truly believe it is essential to take dentistry and dental technology to its next level. The beautiful outcomes we strive for in our daily practice and routines should be made available to as many of our patients as possible—not just the ones who have the financial means to afford comprehensive treatments. As a realist, of course I understand that there will always be

certain limitations to this. However, advancements in digital technology, workflows, and highly esthetic dental materials continue to bring this dream closer to reality. And we should embrace that.

You might ask yourself now if there will be any changes to the previous format and style of QDT. The short answer is no. It is an honor to continue the legacy of such a unique publication with its extraordinary and highly visual content; any major changes would eliminate important parts of QDT DNA. So the only change you will experience over the coming years is a greater variety of international authors. Some of them may even be new to you, but I promise that you will not regret reading and marveling at what they have to share. Finally, and most importantly, I hope to encourage more female dental technicians and clinicians to submit their outstanding (but clearly underrepresented) work to QDT.

Having said all this, it gives me great pleasure to invite you to enjoy this 45th volume of QDT. Stay healthy!

Sincerely yours, Vincent Fehmer vincent.fehmer@unige.ch **kuraray** 



#### Experts Love Kuraray Noritake Porcelain!

"Kuraray Noritake Porcelain. Proven Performance. The results you expect from an industry leader."

- Naoki Hayashi



KATANA Zirconia HTML Structure + CZR Layered #7 Pontic, #8 Screw-retained Implant #7 and #8 are Splinted



"Classification and Indication of Zirconia Restorations according to Layering Design and Thickness." by Naoki Hayashi, RDT, MDT, MDC – 2023 QDT.



#### The extensive product line-up includes:

YML (750 - 1,100 MPa / Translucency 49 - 45%)

UTML (550 MPa / Translucency 50%)

STML (750 MPa / Translucency 49%)

HTML PLUS (1,100 MPa / Translucency 45%)

HT (1,100 MPa / Translucency 43%)

LT (1,100 MPa / Translucency 31%)

#### YML Yttria Multi Layered - The All-In-One Disc Solution



LAYER	FLEXURAL STRENGTH	TRANSLUCENCY
Enamel / 35%	750MPa	49%
Body 1 / 15%	1000MPa	47%
Body 2 / 15%	1100MPa	45%
Body 3 / 35%	1100MPa	45%

The Winning Combination KATANA™ Zirconia and Kuraray Noritake Porcelain





1-844-kuraray (1-844-587-2729) www.katanalabdirect.com









1-800-496-9500 www.zahndental.com

# Enhancing Anterior Esthetics: A Patchwork of Different Types of Geramics



ne of our tasks as dental technicians is to have sufficient knowledge and command of the materials that we use in our practice to achieve an esthetically integrated solution for our patients.

In the case described here, I felt that the most sensible thing to do was to use a variety of different ceramic materials because I had to take three different dental substrates into account.

#### INITIAL SITUATION

The patient complained of pain on the maxillary right lateral incisor. She was also dissatisfied with the appearance of the post-retained crown on the contralateral left lateral incisor. To make matters worse, the central incisors were different shades following endodontic treatment on the left central incisor in the past (Figs 1 to 3). In addition, she had a slight crossbite in the region of her mandibular right canine and first premolar. These minor issues together considerably affected her overall esthetic appearance.

What made this case particularly challenging was the fact that this patient worked as a dental hygienist at her dentist's practice and could spot all the minute details of the soft tissues and teeth that the untrained will not see. Her expectations were therefore high for me and for her boss.

## FIRST APPOINTMENT: DESIGNING THE MOCK-UP

After taking the first impression, I created a wax-up (Fig 4). In the process, I paid particular attention to establishing a symmetric balance and determining the amount of space needed for esthetic improvement in the given clinical circumstances. Therefore, I extended the crown length toward the apical with gingivectomy and modified the labial arch to such an extent that I could correct the crossbite. With the help of a silicone key, these modifications could then be transferred directly to the patient's mouth (Figs 5 and 6).

This kind of mock-up gives me the advantage of being able to immediately assess if the planned modifications will lead to a harmonious integration into the overall appearance. Compared with digital before-and-after visualizations, the mock-up enables the patient to see how the new dental situation looks and feels and also allows her to speak normally.

In this case, the patient was pleased with the new esthetics and feel right away. The slightly extended labial arch of the maxilla suited her, and the optimized proportions of the central incisors with gingivectomy were equally successful. We both agreed to the plan to reconstruct all six maxillary anterior teeth.

The central incisors and canines would be restored with veneers to increase their volume slightly and to redress the slight crossbite on the right side. The maxillary lateral incisors would receive crowns, but the right lateral incisor would first need to be replaced with an implant due to extensive tooth resorption (Fig 7).

\*Dental technician, Dornbirn, Austria

Correspondence to: Carola Wohlgenannt, team@wohlgenannt-zt.com



Figs 1 to 3 Initial situation. Note the different shades of the central incisors in the oblique view.



Fig 4 Wax-up.




Fig 5 Gingivectomy planning.



#### **SECOND APPOINTMENT:** PLACING THE IMMEDIATE **IMPLANT**

The maxillary right lateral incisor was removed surgically, and an implant was placed immediately into the extraction socket (Figs 8 and 9). The implant (Camlog Screw-Line Promote Plus, 3.8 × 13 mm) was placed slightly to the palatal, and the gap between the buccal bone wall and the implant was filled with a heterologous bone mixture (OsteoBiol MP3) and collagen (Fig 10). After careful wound closure, a glass fiber-reinforced Maryland bridge was placed (Fig 11). Just to be prudent, we prepared this bridge prior to the second appointment (Figs 12 to 14).

This choice of treatment provided our patient with some crucial advantages: (1) It eliminated the need for a palatal plate, and (2) it prevented any shifting movement from occurring and therefore protected the newly placed implant effectively.



Figs 8 and 9 The implant is placed slightly to the palatal. Note the gap between the implant and the buccal bone wall.




Fig 10 The gap is filled with a heterologous bone mixture (OsteoBiol MP3).



Fig 11 Glass fiber-reinforced Maryland bridge in situ.



Figs 12 to 14 Laboratory fabrication of the glass fiber-reinforced Maryland bridge with SR Ivocron prior to the second appointment.













18

#### **UNCOVERING THE IMPLANT**

Following a 3-month healing phase, the implant was uncovered and provided with a 4-mm cylindric gingiva former (Figs 15 and 16). In the process, a gingivectomy procedure was performed using an electrotome (Fig 17), and the labial frenulum was moved toward the apical region (Fig 18).

An improvement in the esthetic appearance of the patient could be observed already at this stage.

# PRELIMINARY WORK IN PREPARATION FOR IMPRESSION TAKING

The gingiva former was left in place for 4 weeks before it was replaced with a customized abutment to help establish the emergence profile. The implant was restored with a temporary crown milled from acrylics (Figs 19 and 20). The crown was given a completely nonoccluding design to protect the implant from any unnecessary stresses.

#### **PREPARATION**

The teeth were prepared utilizing a supragingival, minimally invasive approach. This step was performed with the help of the silicone key derived from the wax-up (see Fig 21). Because all the teeth involved in the restoration were being slightly augmented, precious tooth structure could be preserved. In my opinion, however, preparing appropriate interdental spaces was indispensable in this case. I was able to design the contact areas in a targeted fashion and achieve a clean transition between the veneers and the tooth structure by lightly preparing the contact areas in the dorsal area.

Fig 15 Uncovering the implant.

Fig 16 Healing cap.

Fig 17 Gingivectomy using an electrotome.

Fig 18 Cutting the labial frenulum.



The available space was assessed (Fig 21), the teeth were prepared (Figs 22 and 23), and a customized impression post was placed (Fig 24). Subsequently, an impression was taken (Fig 25), and a temporary restoration was placed (Fig 26).



**Fig 21** Space available prior to preparation.



Fig 22 Preparing the two central incisors.

Fig 24 Completed preparation with individualized gingiva former and customized impression post for impression taking.

Fig 23 Completed preparation.





## PLANNING THE PROSTHETIC RESTORATIONS

In this case, the challenge was to achieve a harmonious esthetic appearance of six teeth, four of which contained widely different substrates. It was the dentist's and patient's preference that all the restorations should be made of all-ceramics.

Specifically, the base shade of the two central incisors differed from one another because the left central incisor was devitalized. Another tooth had been restored with a

full-coverage crown on a metal post, and in the region of the right lateral incisor, an implant had been placed following tooth extraction.

The first step in the laboratory was to create a model. I usually use the Giroform model system (Amann Girrbach) to achieve this. This device offers two advantages: First, the rigid baseplate counteracts the contraction of the plaster. Second, the duplication flask supplied with the device enables me to fabricate my refractory dies with accuracy and comparative ease (Figs 27 to 30).





Fig 28 Duplicating the veneer preparations.



Fig 29 Casting the dies using refractory die material.

#### **ALL-CERAMIC SOLUTION**

I now had to figure out a way to establish a common base from which to proceed with my restoration work.

Consequently, I decided to restore the two lateral incisors with copings made of IPS e.max Press lithium disilicate (LT A1, Ivoclar) placed on customized implant abutments milled from zirconia (e.max ZirCAD Prime, BL4). This is how the framework was given strength (Fig 31).

To create ultrathin veneers and natural shade effects, I often opt for IPS Style Ceram (Ivoclar), a metal-ceramic material that comprises a unique blend of leucite, fluoroapatite, and oxyapatite crystals. This combination adds brilliance and vibrancy to the restorations. The shade ranges of IPS e.max Ceram and IPS Style Ceram are coordinated with each other.

After I designed and pressed the two lithium disilicate copings using IPS e.max Press in shade LT A1 (Ivoclar), I built up both lateral incisors (IPS e.max Ceram) in a targeted fashion using the wax-up created before as a reference (Figs 32 and 33).






Fig 31 Completed pressed frameworks made of IPS e.max Press.



It was especially important to support the brightness (value) of the restoration, so I applied a thin layer of Deep Dentin DD BL4 and, directly on top of it, a layer of Power Dentin PD in tooth shade PD A1 (Fig 34). I designed the incisal plate using Incisal I1, and then I contoured the mamelons on top (Fig 35). Next, I applied thin alternate layers of different materials. Then, I applied a small quantity of Impulse Opal Effect OE4 and OE3-OE5 to achieve the bright target shade and to make sure that the value was maintained on the surface (Fig 36).

Having reconstructed the size and shape of the two lateral incisors, I was now ready to focus on the two central incisors and the canines.




Fig 33 Layering and color mapping.



Fig 34 Building up the dentin.



Fig 35 Building up the incisal plate.



**Fig 36** Completing the tooth shape using Effect and Incisal material.

#### REFRACTORY DIES

To create the refractory dies, I took advantage of my flask to duplicate the two incisors and canine preparations and then pour them using Sherarefract refractory die material (Shera). It is essential to thoroughly soak the refractory dies in water before applying the wash (Fig 37). The moist surface prevents the formation of bubbles and prevents the surface from drying prematurely during the subsequent layering procedure.

After soaking the dies, I applied IPS Style Ceram Cervical Transpa Yellow CT, mixed with a small amount of dentin, to the cervical area to make sure that the ceramic material would not delaminate along the margins at the second firing.

I then began the layering process by applying a wash (Fig 38). First I applied a thin layer of IPS Style Ceram Opal Effect OE1 and then I fired this layer. To build up the incisal area, I applied a thin layer of Opal Effect OE1 along the margin and a 50:50 blend of dentin and cervical Transpa material to the cervical area. Then the restoration was fired again.

It was now time to adjust the dentin. In the present case, I first covered the left central incisor with pure Dentin A1 to mask the discolored nonvital tooth. I then continued to build up the dentin, now using a blend of dentin and the matching incisal material (Fig 39). Without this method, there is a risk that the veneers layered with IPS Style Ceram may appear too opaque.

Similar to the IPS e.max crowns, the incisal plate extended from the dentin. In other words, I extended the incisal portion in the direction of the incisal surface but not toward the labial surface. This is important to have enough space to create the internal customizations of the tooth. To complete the buildup, I created the internal structures using mamelon materials (Fig 40). This was followed by alternate layers of different incisal materials in line with the specific requirements of the clinical case (Fig 41). In the present case, this meant using a lot of Opal Effect OE2, OE3, and Transpa Incisal TI.

Bright strips of Opal Effect OE3, mixed with a small amount of OE5, were applied to lend a touch of warmth to the restoration at corrective firing (Fig 42). This way, I achieved the bright Bleach Effect requested by the patient, without incurring the risk of creating a cold, unnatural white shade.






Fig 38 Wash firing using OE 1 including cervical modification firing. Fig 39 Individual dentin layering.



Fig 40 Incisal plate. Covering the mamelon materials with OE3. Fig 41 Completely fired veneering layers.





Figs 43 to 45 Glaze firing and light customizations using stains (Ivocolor).



**Fig 46** Placing veneers on the central incisors.

Next, the existing gold post was covered with opaquer (Fig 47). The lithium disilicate crowns for the lateral incisors were then etched, bonded (Monobond Plus, Ivoclar),

and seated using Multilink Hybrid Abutment dual-curing luting composite. Figures 48 and 49 show the final restorations in place.







Figs 48 and 49 Restorations in situ.

#### CONCLUSION

The esthetic appearance of this discerning patient and professional colleague was successfully restored, enabling her to smile spontaneously and with confidence again (Figs 50 and 51).

Initially, I was worried about how to integrate and harmonize this complex case involving an implant, old crowns,

and a devitalized central incisor. Luckily, my worries proved to be unfounded, and the patient and clinician were happy in the end. Ultimately, it is challenges like this that encourage us to grow in our daily professional lives. Even after many years of experience in the profession, we still come across clinical cases, situations, and valid patient requests that awaken our best qualities and let us use them.



#### Acknowledgment

Time and again, we can see how careful planning is key to any success, and outstanding teamwork is part of this. I would therefore like to extend my thanks to Dr Christian Schult for our long-standing and exceptional collaborative working relationship.





Predictable
Anterior Esthetics
in a Complex Case
of a Young Patient
with a Congenital
Missing Central
Incisor

Arndt Happe, DDS, Dr med dent<sup>1,2,3</sup> Anja Krüssmann<sup>4</sup> Leonie Debring, DMD, MSc<sup>1</sup>